

Infection Rate of *Cryptosporidium parvum* among Diarrheic Children in Isfahan

Hossein Saneian*¹, MD; Omid Yaghini¹, MD; Amene Yaghini², MD;
Mohammad-Reza Modarresi¹, MD, and Mohsen Soroshnia², MD

1. Department of Pediatrics, Isfahan University of Medical Sciences, Isfahan, Iran
2. Isfahan University of Medical Sciences, Isfahan, Iran

Received: May 15, 2009; Final Revision: Feb 19, 2010; Accepted: Apr 21, 2010

Abstract

Objective: Cryptosporidiosis is an important enteric parasitic infection among infants and children in developing countries with significant morbidity and mortality especially among immuno-suppressed individuals. The aim of the present study was to estimate the prevalence of enteric cryptosporidiosis in children presenting with diarrhea in Isfahan.

Methods: This cross-sectional study was conducted from August 2007 to June 2008 in three university hospitals in Isfahan. Children aged 1 month to 10 years presenting with acute or persistent diarrhea were selected consecutively. The oocyst of *C. parvum* was investigated in stool specimens using a modified acid-fast staining method.

Findings: During the study period, 606 children (mean age 42.4±30.0 months, 58.1% female) were recruited. Acute and persistent diarrhea was present in 422 (69.6%) and 184 (30.4%) of the children, respectively. Twenty eight (4.6%) specimens were oocyst positive. The prevalence of *Cryptosporidium* infection was significantly higher in children with persistent diarrhea compared to children with acute diarrhea (12.5% vs 1.2%; $P<0.001$). Most of the infected children were under 5 years of age (89.2%), however, the age difference between infected and non-infected children was not statistically significant. Also, there was no significant difference between infected and non-infected children in gender.

Conclusion: The prevalence of *Cryptosporidiosis* in children presenting with persistent diarrhea is considerable and we suggest routine stool examination for *Cryptosporidium* in this group of children.

Iranian Journal of Pediatrics, Volume 20 (Number 3), September 2010, Pages: 343-347

Key Words: Cryptosporidiosis; *Cryptosporidium parvum*; Diarrhea; Gastroenteritis; Prevalence

Introduction

Diarrhea, including that of parasitic origin, remains one of the most common illnesses

among children and, as reported by the World Health Organization, is one of the major causes of infant and childhood mortality in developing countries^[1]. Intestinal opportunistic parasitic

* Corresponding Author;

Address: Department of Pediatrics, Alzahra Hospital, Soffe Ave, Isfahan, Post Code: 81746-75731, IR Iran

E-mail: saneian@med.mui.ac.ir

infections are important causes of diarrhea which is a serious health problem in tropical regions. *Giardia* spp. and *Cryptosporidium* spp. are common parasitic causes of human diarrhea with the prevalence rate of 1%-3% in the industrialized world and 4%-17% in developing countries^[2]. In developed countries, massive *Cryptosporidium* food borne and waterborne outbreaks have been reported because *Cryptosporidium* oocysts are not killed by conventional disinfectants and chlorination^[3]. In developing countries, *Cryptosporidium* affects mostly children under five years of age and has been reported to be more common in malnourished than in well-nourished children^[4,5].

C. parvum, a protozoan parasite of the phylum *Apicomplexa*, is an enteric pathogen that causes an acute as well as chronic diarrhea in humans. The human host range is broad and includes people with immunodeficiency, children in developing countries, and outbreaks among immuno-competent individuals^[6]. In a healthy adult, infection often results in a self-limited diarrheal illness, however, in children of developing countries, cryptosporidiosis can result in persistent diarrhea leading to malnutrition and developmental delays^[6]. Indeed, *Cryptosporidium* is still an under-diagnosed cause of diarrheal illness in children. Few reports with relatively small sample sizes are available on the prevalence and clinical presentation of *Cryptosporidium* infection in both immuno-compromised and immune-competent children in Iran^[7-13]. The purpose of this study was to determine and compare the prevalence of *C. parvum* infection in children with an acute or persistent diarrhea in Isfahan city, central Iran.

Subjects and Methods

This cross-sectional study was prospectively conducted in three university hospitals in Isfahan from August 2007 to June 2008. Children aged one month to 10 years presenting with acute or persistent diarrhea were visited in each hospital by one of the researchers (pediatrician)

of the study and were included consecutively. Diarrhea was defined as the passage of abnormally liquid or unformed stools at an increased frequency and further defined as acute if lasted <2 weeks and persistent if lasted for 2 to 4 weeks before admission^[14].

Children who had used antiparasitic or antibiotic drugs before admission and also those with known hematologic diseases and immuno-compromised children were excluded from the study. Assuming the average prevalence of *C. parvum* to be 5% according to previous reports^[12], and a margin error of 2% (difference between estimated and reported prevalence) and 95% confidence, a sample size of 456 was calculated for determining the overall prevalence of *C. parvum*.

Fecal samples were collected in sterile plastic bottles and were transported without preservative and arrived in the central laboratory of Isfahan city within 24 hours of collection. The oocyst of *C. parvum* was tested using a modified acid-fast staining method which is a sensitive and specific approach for the identification of *Cryptosporidium* and coccidian oocysts in stools^[15]. In this technique, the oocysts appear as pink to red, spherical to ovoid bodies on a blue or purple background. All samples were evaluated by a single experienced parasitologist. The study protocol was approved by the Ethical Committee of Isfahan University of Medical Sciences and consent was obtained from the parents.

Data analysis was done using SPSS for Windows Version 16.0. The Chi-Square test was used to compare relative frequency of *Cryptosporidium* infection between children with acute and persistent diarrhea. Independent *t*-Test was used to compare the mean age between infected and non-infected children. Statistical significance was set at $P < 0.05$.

Findings

During the study period, 606 children aged one month to 10 years were presented with acute or persistent diarrhea to three university hospitals in Isfahan. There were 254 (41.9%) males and

352 (58.1%) females with the mean age of 42.4 ± 30.0 months. Acute or persistent diarrhea was present in 422 (69.6%) and 184 (30.4%) of the children, respectively.

Total number of *Cryptosporidium* oocyst positive specimens was 28 (4.6%).

Cryptosporidium oocysts were detected in 12.5% (23/184) of the children with persistent compared to 1.2% (5/422) of the children with acute diarrhea; $P < 0.001$. Most (89.2%) of the infected children were under 5 years of age, however, the age difference between the infected and non-infected children was not statistically significant; $P = 0.1$. Also, there was no significant sex difference between the infected and non-infected children; $P = 0.5$ (Table 1).

The age difference between children with acute and persistent diarrhea was not statistically significant (43.9 ± 33.4 vs 38.9 ± 19.6 months, $P = 0.06$) and there was no significant difference between the two sexes (248/174 vs 104/80 female/male ratio, $P = 0.6$).

Discussion

Intestinal parasites are very common in developing countries and *Cryptosporidium* has revealed to be one of the most common parasites. *Cryptosporidium* infests the small intestinal epithelium, thereby, results in an accelerated loss of villous enterocytes, leading to severe villous atrophy and a malabsorptive and secretory diarrhea which is the most pronounced clinical feature of the infection.

Other symptoms include dehydration, weight loss, stomach cramp, or pain, fever, nausea, and vomiting^[6]. The infectious dose is low and

feeding studies have demonstrated that the ingestion of as few as 10-30 oocysts can cause infection in healthy persons^[16]. Infected persons have been reported to shed 10^8 - 10^9 oocysts in a single bowel movement and to excrete oocysts for up to 50 days after cessation of diarrhea^[16,17].

There are discrepancies in the prevalence of cryptosporidiosis in children among surveys in different parts of Iran. The reported prevalence is from 1.5% in Rasht (northern city of Iran) to 21.4% in Shiraz (southern city of Iran)^[7-13]. Our analysis confirmed that cryptosporidiosis is a common infection in children of our community. The difference in the reported prevalence may be attributed to differences in study population (considering age range), diagnostic methods, environmental risk factors (public water supply), time of the study (summer vs winter), nutritional status of the children, and other risk factors^[6,18,19].

Educational level of parents has also been reported as a risk factor^[18]. In other countries, the rates of cryptosporidiosis are 1%-2% in Europe, 0.6%-4.3% in North America, and 3%-10.2% in Asia, Australia, Africa, and Central and South America^[20].

Several methods are available for identification of *Cryptosporidium* oocysts in fecal specimens including modified acid-fast staining which detects oocyst wall, fluorescein-conjugated monoclonal antibody-based detection of oocyst wall antigen, enzyme-linked immunosorbent assay (ELISA) which detects *Cryptosporidium* antigen and most recently polymerase chain reaction (PCR) which detects *Cryptosporidium* DNA^[6]. Modified acid-fast stain of a fecal smear has been the gold standard for detecting *Cryptosporidium* oocysts in stool. This method is commonly used in clinical microbiology laboratories to easily identify

Table 1: Comparison of infected and non-infected children

	Infected n = 28	Non-infected n = 578	P
Age (month)	34.21±16.7	42.81±30.4	0.1*
Female/Male	16(57.1%)/12(42.8%)	336(58.1%)/242(41.8%)	0.5‡
Acute/Persistent	5(17.8%)/23(82.1%)	417(72.1%)/161(27.8%)	< 0.001‡

* t-Test / ‡ Chi-Square Test

Cryptosporidial oocysts. Although the concentration and staining procedures are time-consuming and also require an experienced microscopist to read the slides, it is inexpensive and allows the detection of other parasites (eg, *Isoospora* and *Cyclospora*) at the same time.

However, the sensitivity for the detection of *Cryptosporidium* is substantially better with ELISA formats or immunofluorescent assays^[6].

PCR method is sensitive, but the cost and tedious methodology have limited its use in the clinical microbiology laboratory. Regardless of the method used, it is extremely important that the physician remembers to request Cryptosporidial testing when stool from a child with diarrheal illness is to be examined for ova and parasites^[6].

The results of the present study showed that children with cryptosporidiosis are more likely to present with persistent diarrhea, compared to acute diarrhea. In a report from southeastern Iran, Hamedi and colleagues also found that diarrhea lasted significantly longer in children infected with *Cryptosporidium*^[11]. However, Khalili and colleagues in their study on 171 children found no relationship between type of diarrhea and infection^[10]. Although our study did not show significant age difference between the infected and non-infected children, most of the infected children, as expected, were below 5 years old. Though humans are susceptible at any time in their lives, the prevalence of cryptosporidiosis is higher in younger children mostly in children aged below 5 years^[8,10,11]. The occurrence of high infection rates in younger children might be attributed to their weak immune responses and/or risk of contact with the oocysts. However, more studies are needed to confirm this subject^[12,18].

There are some limitations for the present study. Although the sample size of the study was large enough, sampling was consecutive and not random. However, the study was multicenter and we conducted the study through one year, so it is unlikely that season or selection bias affected our results. Also we applied the modified acid-fast staining and although it is a sensitive and specific approach for the identification of *Cryptosporidium*, ELISA and immunofluorescent assays are more sensitive in

this regard. We recommend evaluating children for important risk factors for infection with *Cryptosporidium*, such as nutritional and socioeconomic status, for future studies.

Conclusion

The prevalence of *Cryptosporidiosis* in children presenting with persistent diarrhea is considerable and we suggest routine stool examination for *Cryptosporidium* in these children for early diagnosis and treatment to prevent possible complications. Also, for control and prevention of this infection, strengthening water quality standards and improving sanitary practices are required.

Acknowledgment

This study was supported by a grant from the Vice Chancellor of Research, Isfahan University of Medical Sciences. We would like to thank Dr Ali Gholamrezaei for editing the manuscript.

Conflict of Interest: None

References

1. Boschi-Pinto C, Velebit L, Shibuya K. Estimating child mortality due to diarrhoea in developing countries. *Bull World Health Organ.* 2008;86(9):710-7.
2. Horman A, Korpela H, Sutinen J, et al. Meta-analysis in assessment of the prevalence and annual incidence of *Giardia* spp. and *Cryptosporidium* spp. infections in humans in the Nordic countries. *Int J Parasitol.* 2004; 34(12):1337-46.
3. DuPont HL, Chappell CL, Sterling CR, et al. The infectivity of *Cryptosporidium parvum* in healthy volunteers. *N Engl J Med.* 1995; 332(13):855-9.
4. Khan WA, Rogers KA, Karim MM, et al. Cryptosporidiosis among Bangladeshi children

- with diarrhea: a prospective, matched, case-control study of clinical features, epidemiology and systemic antibody responses. *Am J Trop Med Hyg.* 2004;71(4):412-9.
5. Steinberg EB, Mendoza CE, Glass R, et al. Prevalence of infection with waterborne pathogens: a seroepidemiologic study in children 6-36 months old in San Juan Sacatepequez, Guatemala. *Am J Trop Med Hyg.* 2004;70(1):83-8.
 6. Huang DB, Chappell C, Okhuysen PC. Cryptosporidiosis in children. *Semin Pediatr Infect Dis.* 2004; 15(4): 253-9.
 7. Khalili B, Shahabi GH, Besharat M, et al. Determining the prevalence of Cryptosporidium and measuring of micronutrients in cryptosporidiosis among children under 5 years in Shahrekord. *J Shaheed Beheshti Univ Med Sci.* 2006;30(3):187-91. (In Persian)
 8. Mohammadi Ghaleh Bin B, Falah S, Asgharzadeh M, et al. Prevalence of Cryptosporidium in children suffering from gastroenteritis in Ardabil hospitals. *J Ardabil Univ Med Sci.* 2006;6(2):176-82. (In Persian)
 9. Berenji F, Zabolnejad N, Kianifar H, et al. Cryptosporidium Infection in Pediatric Patients with Lymphohematopoietic Malignancies. *Iran J Pediatr.* 2007;17(3):247-51.
 10. Khalili B, Shahabi GH, Khayeri S, et al. Prevalence of Cryptosporidium and risk factors related to Cryptosporidiosis in hospitalized children under 5 years of age due to diarrhea (Shahrekord 2005). *Armaghane-Danesh.* 2007;12(3):105-16. (In Persian)
 11. Hamed Y, Safa O, Haidari M. Cryptosporidium infection in diarrheic children in southeastern Iran. *Pediatr Infect Dis J.* 2005;24(1):86-8.
 12. Mirzaei M. Prevalence of Cryptosporidium sp. infection in diarrheic and non-diarrheic humans in Iran. *Korean J Parasitol.* 2007; 45(2):133-7.
 13. Nahrevanian H, Assmar M. Cryptosporidiosis in immunocompromised patients in the Islamic Republic of Iran. *J Microbiol Immunol Infect.* 2008;41(1):74-7.
 14. Ghishan FK. Chronic diarrhea. In: Kliegman RM, Behrman RE, Jenson HB, Stanton BF (eds). *Nelson Textbook of Pediatrics.* 18th ed. New York; Saunders. 2008; Pp:1276-83.
 15. Forbes BA, Sahm DF, Weissfeld AS. Laboratory methods for diagnosis of parasitic infections. In: *Bailey & Scott's Diagnostic Microbiology.* 12th ed. St Louis; Mosby. 2007; Pp: 595-8.
 16. Yoder JS, Beach MJ, Centers for Disease Control and Prevention (CDC). Cryptosporidiosis surveillance -United States, 2003-2005. *MMWR Surveill Summ.* 2007;56(7):1-10.
 17. Smerdon WJ, Nichols T, Chalmers RM, et al. Foot and mouth disease in livestock and reduced cryptosporidiosis in humans, England and Wales. *Emerg Infect Dis.* 2003;9(1):22-8.
 18. Khoshzaban F, Dalimi-Asl A. Epidemiological study on Cryptosporidiosis in diarrheic children. *Daneshwar.* 1998;19(5):21-6. (In Persian)
 19. Roy SL, DeLong SM, Stenzel SA, et al. Risk factors for sporadic cryptosporidiosis among immunocompetent persons in the United States from 1999 to 2001. *J Clin Microbiol* 2004;42(7):2944-51.
 20. Frost FJ, Muller T, Craun GF, et al. Serological analysis of a cryptosporidiosis epidemic. *Int J Epidemiol.* 2000;29(2):376-9.